



Hawley's
Condensed Chemical
Dictionary

THIRTEENTH EDITION

Revised by
Richard J. Lewis, Sr.

BEST AVAILABLE COPY



JOHN WILEY & SONS, INC.

New York • Chichester • Weinheim • Brisbane • Singapore • Toronto

EXHIBIT A

BEST AVAILABLE COPY

This text is printed on acid-free paper. ∞

Copyright © 1997 by John Wiley & Sons, Inc.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning or otherwise, except as permitted under Sections 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 750-4744. Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 605 Third Avenue, New York, NY 10158-0012, (212) 850-6011, fax (212) 850-6008, E-mail: PERMREQ@WILEY.COM.

Library of Congress Cataloging-in-Publication Data

Condensed chemical dictionary.

Hawley's condensed chemical dictionary.—13th ed./revised by
Richard J. Lewis, Sr.

p. cm.
ISBN 0-471-29205-2 (hardcover)

I. Chemistry-Dictionaries. I. Hawley, Gessner Goodrich, 1905-1983.

II. Lewis, Richard J., Sr. III. Title.

QD5.C5 1997

540'.3—dc21

97-35762

CIP

Printed in the United States of America

10 9 8 7 6

alkene. See olefin.

"Alkor" [Atlas]. TM for a synthetic, furan-type resin cement which is acid- and alkali-proof and used as a mortar cement where temperatures do not exceed 380F.

alkoxyaluminum hydrides. (H_nAlOR_n).

A group of reducing agents especially useful in converting epoxides to alcohols. Derived by reaction of aluminum hydride with the corresponding alcohol in tetrahydrofuran.

alkyd resin. A thermosetting coating polymer, chemically similar to polyester resins, conventionally made by condensation and polymerization of a dihydric or polyhydric alcohol (ethylene glycol or glycerol) and a polybasic acid (phthalic anhydride), usually with a drying oil modifier. The process requires heating at 230–250C for up to 12 hours. A new and quite different method utilizes epoxy addition polymerization, in which a mixture of glycidyl esters and organic acid anhydrides is heated with a metallic catalyst at 100C or less for only two to four hours. Cost and energy savings and improved application performance are realized by this process.

Use: Alkyd resins are used as vehicles in exterior house paints, marine paints, and baking enamels. Molded alkyd resins are used for electrical components, distributor caps, encapsulation, and a variety of similar applications.

alkyl. A paraffinic hydrocarbon group which may be derived from an alkane by dropping one hydrogen from the formula. Examples are methyl CH_3^- , ethyl $C_2H_5^-$, propyl $CH_3CH_2CH_2^-$, isopropyl $(CH_3)_2CH_2^-$. Such groups are often represented in formulas by the letter R and have the generic formula C_nH_{2n+1} .
See aryl.

alkylaryl polyethyleneglycol ether.

Use: In surface-active agents.

See isooctylphenoxypolyoxyethylene ethanol for a typical example of this class of compound.

alkylaryl sulfonate. An organic sulfonate of combined aliphatic and aromatic structure, e.g., alkylbenzene sulfonate.

alkylate. (1) A product of alkylation. (2) A term used in the petroleum industry to designate a branched-chain paraffin derived from an isoparaffin and an olefin, e.g., isobutane reacts with ethylene (with catalyst) to form 2,2-dimethylbutane (neohexane). The product is used as a high-octane blending component of aviation and civilian gasolines. (3) In the detergent industry, the term is applied to the reaction product of benzene or its homologs with a long-chain olefin to form an

intermediate, e.g., dodecylbenzene, used in the manufacture of detergents. It also designates a product made from a long-chain normal paraffin that is chlorinated to permit combination with benzene to yield a biodegradable alkylate. The adjectives *hard* and *soft* applied to detergents refer to their ease of decomposition by microorganisms.
See biodegradability; detergent.

alkylation. (1) The introduction of an alkyl radical into an organic molecule. This was one of the early chemical processes used in Germany to furnish intermediates for improved dyes, e.g., dimethylaniline. Other alkylation products are cumene, dodecylbenzene, ethylbenzene, and nonylphenol. (2) A process whereby a high-octane blending component for gasolines is derived from catalytic combination of an isoparaffin and an olefin.
See alkylate (2); neohexane.

alkylbenzene sulfonate. (ABS).

A branched-chain sulfonate type of synthetic detergent, usually a dodecylbenzene or tridecylbenzene sulfonate. Such compounds are known as "hard" detergents because of their resistance to breakdown by microorganisms. They are being replaced by linear sulfonates.

See alkyl sulfonate; linear molecule; detergent; sodium dodecylbenzene sulfonate.

alkyl diaryl phosphate ester. See "Santicizer 141" [Monsanto].

alkyldimethylbenzylammonium chloride.

General name for a quaternary detergent.

See benzalkonium chloride.

alkylene. A phosphated long-chain alcohol.

alkyl fluorophosphate. See diisopropyl fluorophosphate.

alkylolamine. See alkanolamine.

alkyl sulfonate. (linear alkylate sulfonate; LAS). A straight-chain alkylbenzene sulfonate, a detergent specially tailored for biodegradability. The linear alkylates may be normal or iso (branched at the end only), but are C_{10} or longer.
See sodium dodecylbenzene sulfonate.

alkyne. See acetylene hydrocarbon.

Allan-Robinson reaction. Preparation of flavones or isoflavones by condensing *o*-hydroxyaryl ketones with anhydrides of aromatic acids and their sodium salts.

allantoin. (glyoxyldiureide; 5-ureidohydantoin). $C_4H_6N_4O_3$.